

Universal Behavior of Phase Correlations in Non-linear Gravitational Clustering

Peter Watts and Peter Coles

School of Physics & Astronomy, University of Nottingham, University Park, Nottingham NG7 2RD, United Kingdom

`peter.watts@nottingham.ac.uk, peter.coles@nottingham.ac.uk`

Adrian Melott

Department of Physics and Astronomy, University of Kansas, Lawrence, KS 66045

`melott@kusmos.phsx.ukans.edu`

ABSTRACT

The large-scale structure of the Universe is thought to evolve by a process of gravitational amplification from low-amplitude Gaussian noise generated in the early Universe. The later, non-linear stages of gravitation-induced clustering produce phase correlations with well-defined statistical properties. In particular, the distribution of phase differences D between neighboring Fourier modes provides useful insights into the clustering phenomenon. Here we develop an approximate theory for the probability distribution D and test it using a large battery of numerical simulations. We find a remarkable universal form for the distribution which is well described by theoretical arguments.

Subject headings: cosmology: theory — methods: statistical

1. INTRODUCTION

The large-scale structure of the Universe is a complex interconnecting pattern whose structural elements comprise filaments, sheets and clusters of galaxies surrounding large voids. According to standard theories this “cosmic web” develops by a process of gravitational instability from small initial fluctuations in the density of a largely homogeneous early Universe.

The physical description of an inhomogeneous Universe revolves around the dimensionless density contrast, $\delta(\mathbf{x})$, which is obtained from the spatially-varying matter density $\rho(\mathbf{x})$ via $\delta(\mathbf{x}) = [\rho(\mathbf{x}) - \rho_0]/\rho_0$ where ρ_0 is the global mean density. It is useful to expand the density contrast in Fourier series, in which δ is treated as a superposition of plane waves:

$$\delta(\mathbf{x}) = \sum \tilde{\delta}(\mathbf{k}) \exp(i\mathbf{k} \cdot \mathbf{x}). \quad (1)$$

The Fourier transform $\tilde{\delta}(\mathbf{k})$ is complex and therefore possesses both amplitude $C(\mathbf{k}) = |\tilde{\delta}(\mathbf{k})|$ and phase Φ_k where

$$\tilde{\delta}(\mathbf{k}) = C(\mathbf{k}) \exp(i\Phi_k) = A(\mathbf{k}) + iB(\mathbf{k}); \quad (2)$$

the real and imaginary parts are A and B so that $A(\mathbf{k}) = C(\mathbf{k}) \cos(\Phi_k)$ and $B(\mathbf{k}) = C(\mathbf{k}) \sin(\Phi_k)$. We also have $C^2(\mathbf{k}) = A^2(\mathbf{k}) + B^2(\mathbf{k})$ and $\tan \Phi_k = B(\mathbf{k})/A(\mathbf{k})$.

In theories of structure formation involving cosmic inflation (Guth 1981; Guth & Pi 1982; Linde 1982; Albrecht & Steinhardt 1982), the initial fluctuations that seeded the structure formation process form a Gaussian random field (Bardeen et al. 1986) possessing the properties of statistical homogeneity and isotropy. In such fields the real and imaginary parts of $\tilde{\delta}(\mathbf{k})$ are independent Gaussians so that the modulus $C(\mathbf{k})$ has a Rayleigh distribution and the phases Φ_k are uniformly random on the interval $[0, 2\pi]$. Obviously the distribution of δ in position space will also be Gaussian; indeed all finite-dimensional joint probabilities of δ in different locations are multivariate Gaussian in this case (Bardeen et al. 1986).

Even if the primordial density fluctuations were indeed Gaussian, the later stages of gravitational clustering must induce some form of non-linearity. One particular way of looking at this issue is to study the behavior of Fourier modes of the cosmological density field. If the hypothesis of primordial Gaussianity is correct then these modes began with random spatial phases. In the early stages of evolution, the plane-wave components of the density evolve independently like linear waves on the surface of deep water. As the structures grow in mass, they interact with other in non-linear ways, more like waves breaking in shallow water. These mode-mode interactions lead to the generation of coupled phases. While the Fourier phases of a Gaussian field contain no information (they are random), non-linearity generates non-random phases that contain much information about the spatial pattern of the fluctuations but which is ignored entirely in the usual clustering descriptors, such as the power spectrum (see below). There have been a number of attempts to gain quantitative insight into the behavior of phases in gravitational systems. Some studies (Ryden & Gramann 1991; Soda & Suto 1992; Jain & Bertschinger 1998) have concentrated on the evolution of phase shifts for individual modes using perturbation theory and numerical simulations. An alternative approach was adopted by Scherrer, Melott & Shandarin (1991), who developed a practical method for measuring the phase coupling in random fields that could be applied to real data. More recent studies have established connections between phase evolution, clustering dynamics and morphology (Chiang & Coles 2000; Chiang 2001; Chiang, Coles, & Naselsky 2002) and new methods have been developed for visualizing phase information (Coles & Chiang 2000). Connections have also been demonstrated (Watts & Coles 2003) between phase information and alternative measures of non-linear clustering such as the bispectrum (Scoccimarro, Couchman & Frieman 1999; Verde et al. 2000, 2001).

One of the major barriers to the more widespread use of phase-based methods for probing cosmic nonlinearity is the lack of any well-established statistical framework for quantifying the information contained in the Fourier phases. Since phases are angular variables traditional statistical measures of location and dispersion are inappropriate. Moreover, probability distributions for

circular variables have very different properties to those for variables defined on the real line; see the monographs by Mardia (1972) and Fisher (1993) for more detailed discussion. The upshot of this is that it is not yet known whether there are useful standard distributions and limit theorems like those that make the Gaussian distribution so useful and so ubiquitous for statistical analysis. It is this deficiency we address in this letter.

2. STATISTICAL DESCRIPTION

Non-linear clustering is difficult to handle rigorously with analytic methods. Our approach is therefore to develop an approximate theory and test it using numerical experiments. The simulations we shall be comparing to theory are constructed within a finite cubic volume with periodic boundary conditions. The Fourier representation of the clustering pattern they reveal is therefore discrete. In particular phases are defined for wave-vectors on a cubic lattice which, for simplicity, we take to have an integer spacing. In previous analyses of phase coupling phenomena (Chiang & Coles 2000; Chiang 2001; Chiang, Coles, & Naselsky 2002; Coles & Chiang 2000) it has been found useful to introduce a quantity D_k , defined by

$$D_k \equiv \Phi_{k+1} - \Phi_k, \quad (3)$$

which measures the difference in phase of modes with neighboring wave-numbers in one dimension. This has many advantages as a descriptor, not the least of which is that a translation of the origin by x which alters each Φ_k by kx only produces a constant offset in D_k . One can analyze a three-dimensional simulation by extracting a vector (D_1, D_2, D_3) from differences in three orthogonal k -space directions; the result contains information about both the location and structure of features in (x, y, z) space. One can think of D_k as a discrete representation of $d\Phi_k/dk$, the phase gradient. Note that if the two angles Φ_1 and Φ_2 are independent and uniformly random then the difference $\Phi_1 - \Phi_2$ is also uniformly random, so that D_k will be uniformly distributed for Gaussian fields.

When fluctuations are small ($\delta \ll 1$) they evolve linearly: the initial statistics are preserved in this regime because each mode evolves independently. When $\delta \sim 1$, however, mode-coupling terms alter the distributions of both amplitudes and phases. Strictly speaking, therefore, the real and imaginary parts of the Fourier representation of δ in this regime are no longer Gaussian. However the form of the Fourier expansion (1) itself guarantees that, as long as the autocorrelations of δ do not have too large a spatial extent, the Fourier superposition will be approximately Gaussian. It is therefore still a reasonable approximation to take A and B to be Gaussian even for non-linear systems (Fan & Bardeen 1995). More important for our purposes is the fact that, while for Gaussian fields each k -mode is statistically independent, a non-linear field contains mode-mode correlations (Soccimarro, Zaldarriaga & Hui 1999). These induce phase correlations which manifest themselves in departures from uniformity of the distribution of D_k .

We can model the effect by assuming that the two neighboring modes involved in (3) have real and imaginary parts that are Gaussian distributed. For notational ease we suppress the label $\mathbf{k}$ and

so the real parts of each are written A_1 and A_2 and the imaginary parts B_1, B_2 . We assume these can be approximated as Gaussians but we allow them to be cross-correlated. The four variables (A_1, B_1, A_2, B_2) therefore possess a four-dimensional multivariate Gaussian distribution.

If a set of random variables $X_1 \dots X_N$ have a multivariate Gaussian distribution the joint probability density function of the variables has the form

$$\mathcal{P}_N(X_1, \dots, X_N) = \frac{1}{K} \exp\left(-\frac{1}{2} \sum_{i,j} X_i M_{ij}^{-1} X_j\right), \quad (4)$$

where $K = (2\pi)^{N/2}(\det M)^{1/2}$ and the correlation matrix $M_{ij} = \langle X_i X_j \rangle$. For the case we are interested in we can write $\langle A_i^2 \rangle = \langle B_i^2 \rangle = \sigma_i^2$ for $i = 1, 2$. The quantity $\sigma^2(k)$ is related to the power-spectrum, $P(k)$ defined by $P(k) = \langle |\tilde{\delta}(\mathbf{k})|^2 \rangle$. We can also define a quantity α by $\langle A_i A_j \rangle = \langle B_i B_j \rangle \equiv \sigma_i \sigma_j \alpha$ where $\alpha(\mathbf{k})$ parameterizes the cross-correlation of the modes. We also have $\langle A_i B_i \rangle = 0$ (no summation), guaranteeing translation invariance (Chiang & Coles 2000), while $\langle A_i B_j \rangle = -\langle B_i A_j \rangle \equiv \sigma_i \sigma_j \beta$ in a similar fashion to α . Note that α and β may well be equal but we have kept the most general possible form here. These two parameters allow us to construct the required distribution for $\mathcal{P}(A_1, A_2, B_1, B_2)$. We now convert the distribution $\mathcal{P}$ of the vector (A_1, A_2, B_1, B_2) to the distribution $\mathcal{F}$ of the vector $(C_1, \Phi_1, C_2, \Phi_2)$. The result is

$$\mathcal{F} = \frac{C_1 C_2}{4\pi \sigma_1^2 \sigma_2^2 (1 - \chi^2)} \exp(-Q/2), \quad (5)$$

where

$$Q = \frac{\sigma_2^2 C_1^2 + \sigma_1^2 C_2^2 - 2\sigma_1 \sigma_2 C_1 C_2 \chi \cos(\Phi_1 - \Phi_2 + \theta)}{\sigma_1^2 \sigma_2^2 (1 - \chi^2)}. \quad (6)$$

In these expressions we have used $\chi^2 = \alpha^2 + \beta^2$ and $\tan \theta = \alpha/\beta$. Note that if $\chi = 0$ (no mode correlations) this reduces to the product of two Rayleigh distributions for C_1 and C_2 and two uniform distributions of Φ_1 and Φ_2 .

The next step is to construct a conditional distribution of one of the phase angles, say Φ_2 , given specific values for the other three variables, which we designate as c_1, c_2 and ϕ_1 . This can be done straightforwardly using Bayes' theorem. The result is

$$\mathcal{G}(\Phi_2 | c_1, \phi_1, c_2) = \frac{1}{2\pi I_0(\kappa)} \exp[-\kappa \cos(\phi_1 - \Phi_2 + \theta)], \quad (7)$$

where I_0 is a modified Bessel function of order zero and in which $\kappa = c_1 c_2 \chi / \sigma_1 \sigma_2 (1 - \chi^2)$. This conditional distribution shows how the correlation between neighboring phases arises since it depends upon ϕ_1 . It is deficient, however, in that this distribution also depends on specified values for the amplitudes c_1 and c_2 .

A better theory might be obtained by marginalizing over these variables, but the integrals involved are messy. On the other hand the distribution of amplitudes is relatively narrow, peaking around $c_1 = \sigma_1$ and $c_2 = \sigma_2$ and χ is presumably small for quasi-nonlinear stages. Under these

circumstances we expect the distribution of phase differences formed over a large number of pairs to follow the form given above, i.e.

$$P(D) = \frac{1}{2\pi I_0(\kappa)} \exp[-\kappa \cos(D - \mu)]. \quad (8)$$

where μ is the mean angle. This distribution is well-known in the field of circular statistics where it is known as the *von Mises* distribution (Mardia 1972; Fisher 1993). The mean, μ , is controlled by the positions of individual features of the distribution and will consequently vary from sample to sample. The parameter κ , related to χ , describes the level of nonlinearity. When $\kappa \rightarrow 0$ is small the distribution is approximately uniform, while for small κ it takes the form $P(D) \simeq \frac{1}{2\pi}[1 + \kappa \cos(D - \mu)]$ showing that initial departures from random phases manifest themselves as a sinusoidal perturbation of $P(D)$. In the limit $\kappa \rightarrow \infty$ the distribution tends towards a single spike at $\theta = \mu$; this corresponds to a single concentration in position space.

3. NUMERICAL SIMULATIONS

To test our hypothesis with fairly complete coverage of possible parameters, we used a large ensemble of gravitational clustering simulations (Melott & Shandarin 1993). These comprise sets of 128^3 particles and are not large by current standards, but the particular benefit they offer our analysis is a fairly complete coverage of parameter space. There are four realizations of each type of initial conditions, using different pseudo-random number generators to generate the initial phases. Initial power spectra were pure power laws, i.e. $P(k) = Ak^n$ with indices $n = -2, -1, 0$, and 1 . Data is taken every time the scale of clustering doubles, from $k_{\text{nl}} = 64k_f$ to $k_{\text{nl}} = 4k_f$, where k_f is the fundamental mode of the box and k_{nl} is defined by

$$\int_0^{k_{\text{nl}}} P(k) d^3k = 1. \quad (9)$$

The first stage may be significantly compromised by resolution effects; we have simulations with $k_{\text{nl}} = 2k_f$ but these almost certainly suffer from problems connected with the boundary conditions (which are, as usual, periodic).

Analysis of the properties of the phase differences was conducted on each realization of each spectrum and evolutionary phase, i.e. a total of 120 times altogether. We Fourier-transform each stage and extract phases for each wave-vector $\mathbf{k}$. From these we obtain differences D_k in three orthogonal k -space directions from which we form histograms. The mean value μ of each distribution contains information about the specific spatial location of dominant features (Chiang & Coles 2000), which will differ from realization to realization and which also varies with direction. We therefore rotate individual distributions so that they have the same mean value and “stack” the resulting histograms. We also combine all three directional differences into an overall histogram $P(D)$ where $D = (D_1^2 + D_2^2 + D_3^2)^{1/2}/\sqrt{3}$. This approach, together with the large size of the ensemble, produces final histograms with relatively small error bars, as can be seen in Fig 1. Some readers may find it difficult to see the data, as it overlays the theoretical model with high precision.

4. DISCUSSION

The results show extremely good agreement over the range of initial power-spectra and evolutionary stages, although the model does break down at late stages for the case $n = -2$. This is not surprising, given the fact that such spectra have large amounts of power on large scales and phase correlations therefore develop extremely quickly. Even the earliest stage shown of this simulation shows a significantly non-uniform distribution of D . The development of phase correlations with evolutionary stage for each initial spectrum is represented by the increasing deviation from uniformity down each column starting, as expected, with sinusoidal departures.

It is important to check whether deviations in the expansion rate or non power-law initial power spectra would alter these conclusions. Another ensemble of simulations used the power spectrum with CDM shape parameter (Bardeen et al. 1986) $\Gamma = \Omega h = 0.225$. evolved in both an open ($\Omega_{m0} = 0.34$, OCDM), a flat matter-dominated ($\Omega_{m0} = 1$, TCDM), and a flat cosmological-constant background ($\Omega_{m0} = 0.34$, $\Omega_{\lambda 0} = 0.66$, LCDM) All three ensembles were run to an amplitude corresponding to $\sigma_8 = 0.93$. A Hubble Constant $h = 2/3$ was used in the simulation analysis. These N-body runs had three realizations each of 256^3 particles in boxes of side 128 Mpc. The results are shown in figure 2; we found they are also nicely fit with the von Mises distribution. Results are shown also at $z = 1$, which further varies the normalization and background parameters, again with no significant deviation.

The excellent match of the distribution (8) to the results of detailed numerical simulations may appear surprising given the very approximate nature of its derivation. But its validity is reinforced by the fact that it is the maximum entropy distribution on a circle for a fixed mean μ and fixed circular dispersion (Mardia 1972). It should really therefore be regarded as the circular equivalent of a Gaussian distribution, which has maximal entropy for fixed mean and variance on the real line.

The universal behavior we have demonstrated will allow us in future to discriminate between gravitationally-induced mode-coupling and other forms, such as that induced by peculiar motions (Melott et al. 1998). It should be pointed out, of course, that in a realistic application to a survey of galaxies, further coupling between the Fourier modes would arise due to the geometry and selection function of the survey. However, this need not be problematic. If one knew the window function sufficiently accurately its Fourier transform could be computed and a correction applied to the complex Fourier components themselves.

This work was supported by PPARC grant PPA/G/S/1999/00660; ALM gratefully acknowledges the support of NSF through grant AST-0070702, and computing support from the National Center for Supercomputing Applications.

REFERENCES

- Albrecht, A., and Steinhardt, P. J. 1982, Phys. Rev. Lett., 48, 1220
- Bardeen, J. M., Bond, J. R., Kaiser, N., and Szalay, A. S. 1986, ApJ, 304, 15
- Chiang, L.-Y. 2001, MNRAS, 325 405
- Chiang, L.-Y., and Coles, P. 2000, MNRAS, 311, 809
- Chiang, L.-Y., Coles, P., and Naselsky, P. D. 2002, MNRAS, 337, 488
- Coles, P., and Chiang, L.-Y. 2000, Nature, 406, 376
- Fan, Z., and Bardeen, J. M. 1995, Phys. Rev. D., 51, 6714
- Fisher, N. I. 1993, Statistical Analysis of Circular Data (Cambridge: Cambridge University Press)
- Guth, A. H. 1981, Phys. Rev. D., 23, 347
- Guth, A. H. and Pi, S.-Y. 1982, Phys. Rev. Lett., 49, 1110
- Jain, B., and Bertschinger, E. 1998, ApJ, 509, 517
- Linde, A. 1982, Phys. Lett. , B108, 389
- Mardia, K. V. 1972, Statistics of Directional Data (London: Academic Press)
- Melott, A. L., Coles, P., Feldman, H. A., and Wilhite, B. 1998, ApJ, 496, L85
- Melott, A. L., and Shandarin, S. F. 1993, ApJ, 410, 469
- Ryden, B. S., and Gramann, M. 1991, ApJ, 383, L33
- Scherrer, R. J., Melott, A. L., and Shandarin, S. F. 1991, ApJ, 377, 29
- Scoccimarro, R., Couchman, H. M. P., and Frieman, J. A. 1999, ApJ, 517, 531
- Scoccimarro, R., Zaldarriaga, M., and Hui, L. 1999, ApJ, 527, 1
- Soda, J., and Suto, Y. 1992, ApJ, 396, 379
- Verde, L., Jimenez, R., Kamionkowski, M., and Matarrese, S. 2001, MNRAS, 325, 412
- Verde, L., Wang, L., Heavens, A. F., and Kamionkowski, M. 2000, MNRAS, 313, 141
- Watts, P. I. R., and Coles, P. 2003, MNRAS, 338, 806




Fig. 1.— Histograms $P(D)$ of phase differences D are shown in grey along with the analytical model (8) for four different initial spectra as functions of epoch. Close agreement makes it difficult to discriminate between data and theory.




Fig. 2.— Histograms $P(D)$ of phase differences D are shown in grey along with the analytical model (8) for three different CDM models as functions of epoch.